REFERENCES

1 Azeredo L. A. I., M. B. Lima, R. R. R. Coelho, D. M. G. Freire. (2006). Thermophilic protease production by *Streptomyces* sp. 594 in submerged and solid-state fermentations using feather meal. Journal of Applied Microbiology 100:4.

2 Barrett.A.J.,Rawlinga N.D. ,Brion E.A.,(2001).The MEROPS database as protease information system. Journal of Structural Biology; 14:95-102.

3 Bockle B, Galunsky B,Muller(1995). Characterization of a keratinolytic serine proteinase from Streptomyces pactum DMS 40530. Applied Environmental Microbiology 61: 3705-3710.

4 Brandellli, A and A. Riffel,(2006). Keratinolytic bacteria isolated from feather.Brazilian journal of Microbiology; 37:395-399.

5 Casarin F.,Cladera-Olivera F. Brandell.A. (2008).Use of poultry byproduct for production of keratinolytic Enzymes .Food and Bioprocess Technology;3: 301-305.

6 Coward –Kelly G,Frank K,Agbogbo A. , Mark T.,(2006).Holtzapple lime treatment of keratinous materials for the generation of highly digestible animal feed.Animal hair Bioresource Technology; 97: 1344-1352.

7 Chitte R. R., V. K. Nalawade and S. De (1998). Keratinolytic activity from the broth of a feather degrading thermophilic Streptomyces thermoviolaceus strain *SD8*. Letters in applied microbiology; 28:131 - 136.

8 Cheng S. W., H. M. Hu and S. W. Sen. (1995). Production and characterization of a keratinase of feather degrading Bacillus licheniformis PWD – 1. Bioscience biotechnology and biochemistry. 59: 2239-2243.

9 Dalev P., Friedrichet et al (1999) .An enzyme-alkaline hydrolysis of feather keratin for obtaining a protein concentrate for fodder,Biotechnology Letters; 12: 71-72.

10 George, S.Raju, V. Krishna, M.R.V,Subramaniun, T. E.and Jayaraman, K.(1995).Production of protease by Bacillus amiloliquefacience in solid state fermentation and its application in the unhairing of hides and skin .Process Biochemistry;30:457-462.

11 Gradisar H,Friedrich J,Krizaj I, Jerala R. (2005) Similarities and specificities of fungal keratinolytic proteases : comparison of keratinase of Paecilomyces Marquandii and Doratomyces microspores to some known proteases .Applied Environmental Microbiology;71: 3420-3426.

12 Holtz,J.D. Williams and Wilkins ,Baltimore.(1997) Bergey's Manual of Determinative Bacteriology .

13 Ichida J.M., L Krizova, C.A. Lefevre, H.M .Keener, D.L. Elwell and E.B. Burtt. (2001). Bacterial inoculum enhaces keratine degradation and biofilm formation in poultry compost. J. Microbiol. *Methods* .47: 99-208.

14 Jadhav A.R. and A. M. Deshmukh. (2005). Keratinolytric activity of *Streptomycete fradiae* isolated from Rankala Lake water Kolhapur Maharashtra. J. Aqua. Biol;.20 :1-4.

15 Jahan H. and R. K. S. Kushwaha. (1999). Human hair degradation by two strains of *Chrysosporium* Tropicum. J. Microb. World. 1(1):61-66.

16 Joshi.S., Tejaswini M.M.,Revati N,Sridevi R., Roma D. (2007). Isolation , Identification and characterization of a Feather Degrading Bacterium .International Journal of Poultry Science; 6: 689-693.

17 Kane, W. D. (1966). A new genus of Actinoplanaceae, Pilimelia, with a description of two Pilimelia terevasa and Pilimelia anulata.J Elisha Mitchell Sci Soc 82,220-230.

18 Kushwaha R. K. S. (1983). The in vitro degradation of peacock feathers by some fungi: *Mykosen* 26:324–326.

19 Latshaw J.D., Musharaf N.,Returm R. (1994). Processing of feather meal to maximize its nutritional value for Poultry,Anim.Feed Sci, Technol;47:.179-188,

20 Letourneau F., V. Soussotte, P. Bressollier, P. Branland and B. Verneuil. (1997). Keratinolytic activity of *Streptomyces sp. S. K1-02*: a new isolated strain. Letters in applied microbiology; 26:77 - 80.

21. Lin, X.; C.G.Lee,; E.S. Casaleand J.C.H. Shih, (1995). Purification and characterization of a keratinase from a feather-degrading Bacillus licheniformis strain. Applied and Environmental Microbiology; 58: 3271-3275.

22 Lin, X, G.D. Inglis; L.J. Yanke and K.J. Cheng. (1999). Selection and characterization of feather degrading bacteria from canola meal compost. Journal of Industrial Microbiology and Biotechnology; 23:149-153.

23 Lowry O.H., N.J. Rosebrough, A.L. Farr and R.J. Randall. (1951). Protein measurement with the Folin phenol reagent. J. Biol. Chem. 193: 265-275.

24 Lyndal M.Pettett and D. Ipek Kurtboke. (2004). Development of an environmentally friendly biofertilizer with keratin degrading and antibiotic producing actinomycetes: Actinoycetologica. 18(2)

25 Minghai Han,Wei Luo,Quiya Gu and Xiaobin Yu., (2011).Isolation and characterization of keratinolytic Protease from a feather degrading bacterium Pseudomonos aeruginosa C II .African Journal of Microbiology Research;l.6:.2211-2221.

26 Mohamed A.Hassan,Bakry M.Haroun, volume (2014).Production and characterization of keratinolytic Protease from New wool degrading Bacillus species isolated from Egyptian Ecosystem.Biomedical Research International.

27 Mukhopadyay, R. P., and A. L. Chandra.(1990). Keratinase of a *streptomycete*. Indian J. Exp. Biol; 28:575–577.

28 Mukhopadyay, R. P., and A. L. Chandra. (1993). Protease of keratinolytic *Streptomycete* to unhair goat skin. Indian J. Exp. Biol; 31:557–558.

29 Noval J.J., W. Nickerson. (1959). Decomposition of native keratin by *Streptomyces fradiae*. J. Bacteriol; 77: 251-263.

30 Onifade A.A.,Al-Sane N.A.,Al-Musallam A.A.,Al-Zarbans., (1998)A review , Potential for biotechnological applications of keratin degrading microorganisms and their enzymes for nutritional improvement of feathers and other keratins of livestock feed resources .Bioresource Technology;66:1-11,.

31 - Papadopoulos M.C.,El Boushy A.R., Ketelaars E.H., (1985). Effect of different Processing conditions on amino acid digestibility of feather meal determined by chicken assay.Poultry .Sci; 64: 172-174.

32- Periasamy Anbu, Hilda A. and S. C. B. Gopinath. (2004). Keratinophilic fungi of poultry farm and feather dumping soil in Tamil Nadu India: Mycopathologia; 158:303–309.

33 Pushpalata Kainoor,G.R.Naik.(2010).Production and Characterization of feather degrading Keratinase from Bacillus Spp.JB 99. Indian Journal of Biotechnology ;9: 384-390.

34 Rajak, R. C., H. K. Malviya, H. Deshpande, and S. K. Hasija. (1992). Keratinolysis by *Absidia cylindrospora* and *Rhizomucor pusillus*: biochemical proof. Mycopathologia 118:109–114.

35 Rao,M.B.; Tanksale,A.M.; Ghatge, M.S.and Deshpande, V.V. (1998). Molecular and biotechnological aspects of microbial proteases. Microbiology and Molecular Biology Reviews; 62: 597-635.

36 Revathi K, Shafali Singh ,Mohd.Azeem Khan, Suneetha V.(2013).A Potential strain of Keratinolytic bacteria VIT RSAS2 from Katpadi and its Pharmacolocgical Benefits.Intrnational Journal of Pharmacological Science;.20:2

37 Riffel, A.; Lucas, F.S.; Heeb, P. and Brandelli, A. (2003). Characterization of a new keratinolytic bacterium that completely degrades native feather keratin. Archives of Microbiology; 17:258-265.

38 Salminen E.and Rintala J., (2002)Semicontinuous anaerobic digestion of solid poultry slaughter house waste, effect of hydraulic retention time and loading,Water Research; 36:3175-3182,.

39 Salvam K and Vishnupriya,B . (2012)Biochemical and Molecular characterization of Microbial keratinase and its remarkable application.,International journal of Pharmaceutical and Biological Archives;3:267-275.

40 Sarita Agrahari and Neeraj Wadhwa .(2010).Degradation of chicken feather a Poultry waste product by a keratinolytic Bacterium isolated from Dumping site at Ghazipur Poultry Processing plant.,International Journal of Poultry Science;9:482-489.

41 Savitha G. Joshi,M.M.Tejaswini,N.Revati,R.Sridevi and D.Roma ., (2007)Isolation ,Identification and characterization of a feather degrading bacterium .International Journal of Poultry Science; 6:689-693.

42 Sangeeta Lal , R. C. Rajak, and S. K. Hasija. (1999). In vitro degradation of keratin by two species of Bacillus: Journal of.General Applied. Microbiology; 45: 283 – 287.

43 Sangali, S. and Brandelli, A. (2000). Feather keratin hydrolysis by a *Vibrio* sp. strain kr2. Journal of Applied Microbiology; 89: 735-743.

44 Suneeta V,V.V. Lakshimi. (2005). Optimization of parameters for fermentative production of Keratinase by feather degrading microorganisms. Journal of Microbial World. 7(1):106 - 115.

45 Scott J. A. and W. A. Untereiner. (2004). Determination of keratin degradation by fungi using keratin azure. Medical mycology; 42: 239-246.

46 Takami H.,N.Satoshi,R.Aono,and K.Horikoshi. (1992).Degradation of human hair by a thermostable alkaline protease from alkaline Bacillus sp.no.AH 101.Biosc.Biotechnol. Biochem;56:1667-1669.

47 Tamilmani P., Umamaheshwari A., Vinayagam A.and B.Prakash , (2008).Production of an Extracellular Feather degrading Enzyme by Bacillus licheniformis isolated from Poultry Farm soil in Namakkal District .International Journal of Poultry Science; 7 :184-188,.

48 Thanikaivelan P., Rao J. R., Nair B. U. Amasam I T. (2004). Progress and recent trends in biotechnological methods for leather processing. Trends in Biotechnology; 22: 181-188.

49 Wawrzkiewicz ,K.,T.Wolski, and J.Lobarzewski (1991).Screening the keratinolytic activity of dermatophytes in vitro.Mycopathologia; 114:1-8.

50 Williams C.M., C.S. Richter, J.R.J.M Mackenzie and J.C.H. Shih. (1990). Isolation, identification and characterization of a fearther-degrading bacterium. Applied Evironmental Microbiology; 56: 1509-1515.

51 Williams, C.M.; Lee, C.G.; Garlich, J.D. and Shih, J.C.H. (1991). Evaluation of a bacterial feather fermentation product, feather-lysate, as a feed protein. Poultry Science; 70: 85-94.

52 Wang J. J. and J. C. H. Shih., (1999). Fermentative production of keratinase from *Bacillus licheniformis PWD – 1* and a recombinant *B. subtilis* FDB-29. J of Industrial microbiology and biotechnology; 22: 608-616.